

Cryopreservation of gametes and gonadal tissues for porcine genetic resources

Kazuhiro Kikuchi^{1,2,*}, Hiroyuki Kaneko¹, Michiko Nakai¹, Tamas Somfai³,
Naomi Kashiwazaki⁴, Takashi Nagai⁵

¹Institute of Agrobiological Sciences, National Agriculture and Food Research Organization (NARO), Tsukuba 305-8602, Japan,

²The United Graduate School of Veterinary Science, Yamaguchi University, Yamaguchi, 753-8511, Japan,

³Institute of Livestock and Grassland Science, NARO, Tsukuba 305-0901, Japan,

⁴Graduate School of Veterinary Science, Azabu University, Sagamihara, 252-5201, Japan,

⁵Past address: Headquarter, NARO, Tsukuba, 305-8517, Japan

*Corresponding author: kiku@affrc.go.jp

Keywords: pig, cryopreservation, vitrification, oocyte, gonadal tissue, gene bank

Abstract

The conservation or preservation of mammalian genetic resources, especially farm animals, has been conducted under *in situ* conditions by maintaining living individuals as "livestock". However, systems for laboratory *in vitro* embryo production using gametes such as spermatozoa and oocytes are now available and *in vitro* culture for embryos viable to offspring after transfer to recipients, as an *ex situ* preservation method for mammalian genetic resources. In pigs, freezing of sperm is the most reliable and well-established method for this purpose. On the other hand, cryopreservation of female gametes (oocytes) and gonadal (testicular and ovarian) tissues usually by vitrification has been conducted; however, resulted in very low efficacies. Recently, our laboratory conducted some research themes related to these issues. We have been focusing on advances using porcine *in vitro* embryo production (IVP) systems and xenografting into host immunodeficient mice for piglet generation. In this symposium, we introduce recent progress on the vitrification of porcine immature oocytes and gonadal tissues followed by their IVP and xenografting to produce gametes.

Introduction

The main type of genetic material for routine cryopreservation by gene banking is spermatozoa, so called semen cryobanking. Banking of other materials such as

oocytes or early embryos has been considered challenging. However, the utilization of female gametes and gonadal tissues has been considered very important. Some attempts at vitrification of tissues have been made, and for this purpose, *in vitro* embryo production (IVP) and its related technologies are essential.

The basic IVP procedure for piglet production has been of fundamental importance in studies of embryo freezing/vitrification at the pronucleus (Somfai et al. 2009), the 4- to 8-cell (Nagashima et al., 2007), the morula (Maehara et al., 2012), and the blastocyst (Mito et al., 2015) stages. Intracytoplasmic sperm injection (ICSI)-related technologies (Nakai et al., 2003, 2009, 2010, Kaneko et al., 2013) are also fundamental for the utilization sperm without self-penetration ability into oocytes.

Cryopreservation of unfertilized oocytes

Freezing of spermatozoa is one of the basic approaches for preservation of genetic resources. Oocyte cryopreservation is a basic strategy for gene banking of female germplasm; for pigs. However, this technology is very difficult and still considered a challenge.

We have tested the possibility of vitrification for immature (at the germinal vesicle stage; cumulus oocyte complexes) and mature (at the metaphase-II stage) oocytes before fertilization (Somfai et al., 2012). The survival rate for mature oocytes was relatively high; however, a high

proportion (about 70%) of vitrified immature oocytes was able to complete meiotic maturation and undergo normal fertilization and embryo development. Using cryopreserved immature oocytes, we had focused on piglet production after *in vitro* maturation/fertilization (IVM/IVF) and embryo transfer, and finally obtained piglets (Somfai et al., 2014).

This application has been confirmed using immature oocytes from commercial Western pigs, which seems to be also efficient reproduction system in native species especially, in Asian countries. We have conducted under the project by Science and Technology Research Partnership for Sustainable Development (SATREPS) for Vietnamese native pigs (https://www.jst.go.jp/global/english/kadai/h2604_vietnam.html) and found equal efficacy in cryopreservation efficacies (Somfai et al., unpublished data).

Cryopreservation of gonadal tissues

1) Testicular tissue

One approach for inducing spermatogenesis in isolated testicular tissues is ectopic (into another site in the body) grafting into immunodeficient host animals (xenografting). Ectopic xenografting has been reported for hamsters (Schlatt et al., 2002), goats (Honaramooz et al., 2002), rhesus macaques (Honaramooz et al., 2004), sheep (Zeng et al., 2006), cats (Snedaker et al., 2004), and pigs (Nakai et al., 2009, Nakai et al., 2010, Honaramooz et al., 2002, Zeng et al., 2006, 2007, Honaramooz et al., 2008, Kaneko et al., 2008). It is preferable to graft testicular tissue under the skin of the back of commercially available severe combined immunodeficiency mice or nude mice.

A series of studies has been conducted in our laboratory (Nakai et al., 2009, Nakai et al., 2010, Kaneko et al., 2008, Kikuchi et al., 2011) to evaluate whether boar spermatogonia can develop into sperm in testicular tissues grafted into immunodeficient nude mice, and whether live piglets can be produced from this sperm by ICSI into oocytes. Elongated spermatids and spermatozoa

were obtained after 4 months. Just after collection, some spermatozoa showed only faint motility for a short period; however, they could not penetrate into IVM oocytes, meaning that we unfortunately could not apply these sperm for IVP and have to select ICSI method to obtain fertilized oocytes. After culture, some of them developed into blastocysts; which were similar to the oocytes from prepubertal gilts after IVM/IVF (Kikuchi et al., 2002, Nakai et al., 2009]. When oocytes at the pronuclear stage after ICSI were transferred to oviducts of estrous-synchronized recipients, we were able to obtain both male and female piglets (Nakai et al., 2010), which showed normal reproductive abilities when developed to the adulthood (Kaneko et al., 2012).

For more advanced utilization of this technique, we have investigated the possibility of vitrification of testicular tissue fragments before xenografting. This method enables long term storage in liquid nitrogen of the tissue and the production of sperm whenever the need arises. After a certain period (120 days), sperm were first recovered from the grafts. The sperm recovery rate increased with time after grafting from 180 to 350 days. Sperm after the recovery from the tissues were applied for ICSI, and the oocytes were transferred to recipients. We could obtain live male and female piglets (Kaneko et al., 2013). Both the male and female pigs showed normal reproductive abilities (Kaneko et al., 2014).

2) Ovarian tissues

Primordial follicles act as stores for ovarian follicles and can be potential sources of oocytes. To utilize these resources, ovarian tissue grafting is a possible method for growing or maturing such primordial oocytes in the primordial follicles of large mammals. In humans and primates, grafting of ovarian tissues to another site in the body (autografting) has been used successfully to produce viable offspring (Lee et al., 2004, Donnez et al., 2004). Xenografting is more advantageous for the conservation and multiplication of domestic or endangered animals. Some studies have been conducted in cat (Gosden et al., 1994), sheep (Gosden et al., 1994), mice (Snow et al.,

2002), humans, (Oktay et al., 1998, Weissman et al., 1999, Kim et al., 2002, Gook et al., 2003), dogs (Metcalf et al., 2001), monkeys (Candy et al., 1995), cattle (Senbon et al., 2003), pigs (Kaneko et al., 2003), tammar wallabies (Mattiske et al., 2002), and common wombats (Cleary et al., 2003, 2004).

Our studies using pigs were the first to demonstrate clearly that such oocytes were competent in terms of fertilization ability (Kaneko et al., 2003) and had limited potential for development into embryos (Kaneko et al., 2006, Kikuchi et al., 2006, 2011); however, we did not have not yet obtain any piglet. From the viewpoint of genetic resource conservation, cryopreservation of ovarian tissues before xenografting may be important. We are trying to confirm this possibility and reported the fertilizing ability (Kikuchi et al., 2010), but its efficacy still remains quite low.

Conclusion and Perspective

Although pigs are important as a source of meat, they are also expected to have considerable potential as models or donors in human regenerative medicine. For example, the possibility of generating the human pancreas from human induced pluripotent stem cells using genetically modified pigs has been suggested. For these purposes, the conservation of porcine genetic resource is very important. We are now encouraged to establish the conservation technologies in pigs.

Acknowledgements

Studies in this review article were supported in part by a grant-in-aid for Scientific Research from the Japanese Society for the Promotion of Science (JSPS) awarded to K. Kikuchi (26480462) and H. Kaneko (26292171), and also supported by Science and Technology Research Partnership for Sustainable Development (SATREPS) from Japan Science and Technology Agency (JST)/Japan International Cooperation Agency (JICA) awarded to K. Kikuchi and T. Somfai.

References

- Candy, C.J., Wood, J. and Whittingham, D.G. 1995. Follicular development in cryopreserved marmoset ovarian tissue after transplantation. *Hum Reprod.* 10: 2334-2338.
- Cleary, M., Paris, M.C., Shaw, J., Jenkin, G. and Trounson, A. 2003. Effect of ovariectomy and graft position on cryopreserved common wombat (*Vombatus ursinus*) ovarian tissue following xenografting to nude mice. *Reprod Fertil Dev.* 15: 333-342.
- Cleary, M., Shaw, J.M., Jenkin, G. and Trounson, A.O. 2004. Influence of hormone environment and donor age on cryopreserved common wombat (*Vombatus ursinus*) ovarian tissue xenografted into nude mice. *Reprod Fertil Dev.* 16: 699-707.
- Donnez, J., Dolmans, M.M., Demylle, D., Jadoul, P., Pirard, C., Squifflet, J., Martinez-Madrid, B. and van Langendonck, A. 2004. Live birth after orthotopic transplantation of cryo-preserved ovarian tissue. *Lancet.* 364: 1405-1410.
- Gook, D.A., Edgar, D.H., Borg, J., Archer, J., Lutjen, P.J. and McBain, J.C. 2003. Oocyte maturation, follicle rupture and luteinization in human cryopreserved ovarian tissue following xenografting. *Hum Reprod.* 18: 1772-1781.
- Gosden, R.G., Boulton, M.I., Grant, K. and Webb, R. 1994. Follicular development from ovarian xenografts in SCID mice. *J Reprod Fertil.* 101: 619-623.
- Honaramooz, A., Snedaker, A., Boiani, M., Scholer, H., Dobrinski, I. and Schlatt, S. 2002. Sperm from neonatal mammalian testes grafted in mice. *Nature.* 418:778-781.
- Honaramooz, A., Li, M.W., Penedo, M.C.T, Meyers, S. and Dobrinski, I. 2004. Accelerated maturation of primate testis by xenografting into mice. *Biol Reprod.* 70: 1500-1503.
- Honaramooz, A., Cui, X.S., Kim, N.H. and Dobrinski, I. 2008. Porcine embryos produced after intracytoplasmic sperm injection using xenogeneic pig sperm from neonatal testis tissue grafted in mice. *Reprod Fertil Dev.* 20: 802-807.

- Kaneko, H., Kikuchi, K., Noguchi, J., Hosoe, M. and Akita, T. 2003. Maturation and fertilization of porcine oocytes from primordial follicles by a combination of xenografting and in vitro culture. *Biol Reprod.* 69: 1488-1493.
- Kaneko, H., Kikuchi, K., Nakai, M. and Noguchi, J. 2008. Endocrine status and development of porcine testicular tissues in host mice. *J Reprod Dev.* 54: 480-485.
- Kaneko, H., Kikuchi, K., Noguchi, J., Ozawa, M., Ohnuma, K., Maedomari, N. and Kashiwazaki, N. 2006. Effects of gonadotropin treatments on meiotic and developmental competence of oocytes in porcine primordial follicles following xenografting to nude mice. *Reproduction.* 131: 279-288.
- Kaneko, H., Kikuchi, K., Nakai, M., Tanihara, F., Noguchi J., Noguchi, M., Ito, J. and Kashiwazaki, N. 2012. Normal reproductive development of offspring derived by intracytoplasmic injection of porcine sperm grown in host mice. *Theriogenology.* 78: 898-906.
- Kaneko, H., Kikuchi, K., Nakai, M., Somfai, T., Noguchi, J., Tanihara, F., Ito, J. and Kashiwazaki, N. 2013. Generation of live piglets for the first time using sperm retrieved from immature testicular tissue cryopreserved and grafted into nude mice. *PLoS One.* 8: e70989.
- Kaneko, H., Kikuchi, K., Tanihara, F., Noguchi, J., Nakai, M., Ito, J. and Kashiwazaki, N. 2014. Normal reproductive development of pigs produced using sperm retrieved from immature testicular tissue cryo-preserved and grafted into nude mice. *Theriogenology.* 82: 325-331.
- Kikuchi, K., Onishi, A., Kashiwazaki, N., Iwamoto, M., Noguchi, J., Kaneko, H., Akita, T. and Nagai, N. 2002. Successful piglet production after transfer of blastocysts produced by a modified *in vitro* system. *Biol Reprod.* 66: 1033-1041.
- Kikuchi, K., Kaneko, H., Nakai, M., Noguchi, J., Ozawa, M., Ohnuma, K. and Kashiwazaki, N. 2006. *In vitro* and *in vivo* developmental ability of oocytes derived from porcine primordial follicles xenografted into nude mice. *J Reprod Dev.* 52: 51-57.
- Kikuchi, K., Kashiwazaki, N., Nakai, M., Noguchi, J., Ito, J. and Kaneko, H. 2010. In vitro maturation and fertilization of oocytes from ovarian tissues cryopreserved and xenografted into nude mice (Abstract). *Reprod Fertil Dev.* 22: 206-207.
- Kikuchi, K., Nakai, M., Kashiwazaki, N. and Kaneko, H. 2011. Xenografting of gonadal tissues into mice as a possible method for conservation and utilization of porcine genetic resources. *Anim Sci J.* 82: 495-503
- Kim, S.S., Soules, M.R. and Battaglia D.E. 2002. Follicular development, ovulation, and corpus luteum formation in cryopreserved human ovarian tissue after xenotransplantation. *Fertil Steril.* 78: 77-82.
- Lee, D.M., Yeoman, R.R., Battaglia, D.E., Stouffer, R.L., Zelinski-Wooten M.B., Fanton, J.W. and Wolf, D.P. 2004. Live birth after ovarian tissue transplant. *Nature.* 428: 137-138.
- Maehara, M., Matsunari, H., Honda, K., Nakano, K., Takeuchi, Y., Kanai, K., Matsuda, T., Matsumura, Y., Hagiwara, Y., Sasayama, N., Shirasu, A., Takahashi, M., Watanabe, M., Umeyama, K., Hanazono, Y. and Nagashima, H. 2012. Hollow fiber vitrification provides a novel method for cryopreserving *in vitro* maturation/fertilization-derived porcine embryos. *Biol Reprod.* 87: 133.
- Mattiske, D., Shaw, G. and Shaw, J.M. 2002. Influence of donor age on development of gonadal tissue from pouch young of the tammar wallaby, *Macropus eugenii*, after cryopreservation and xenografting into mice. *Reproduction.* 123: 143-153.
- Metcalfe, S.S., Shaw, J.M. and Gunn I.M. 2001. Xenografting of canine ovarian tissue to ovariectomized severe combined immunodeficient (SCID) mice. *J Reprod Fertil Suppl.* 57: 323-329.
- Mito, T., Yoshioka, K., Noguchi, M., Yamashita, S., Misumi, K., Hoshi, T. and Hoshi, H. 2015. Birth of piglets from in vitro-produced porcine blastocysts vitrified and warmed in a chemically defined medium. *Theriogenology.* 84: 1314-1320.

- Nagashima, H., Hiruma, K., Saito, H., Tomii, R., Ueno, S., Nakayama, N., Matsunari, H. and Kurome, M. 2007. Production of live piglets following cryopreservation of embryos derived from *in vitro*-matured oocytes. *Biol Reprod.* 76: 900-905.
- Nakai, M., Kashiwazaki, K., Takizawa, A., Hayashi, Y., Nakatsukasa, E., Fuchimoto, D., Noguchi, J., Kaneko, H., Shino, M. and Kikuchi, K. 2003. Viable piglets generated from porcine oocytes matured *in vitro* and fertilized by intracytoplasmic sperm head injection. *Biol Reprod.* 68: 1003-1008.
- Nakai, M., Kaneko, H., Somfai, T., Maedomari, N., Ozawa, M., Noguchi, J., Kashiwazaki, K. and Kikuchi, K. 2009. Generation of porcine diploid blastocysts after injection of spermatozoa grown in nude mice. *Theriogenology.* 72: 2-9.
- Nakai, M., Kaneko, H., Somfai, T., Maedomari, N., Ozawa, M., Noguchi, J., Ito, J., Kashiwazaki, N. and Kikuchi, K. 2010. Production of viable piglets for the first time using sperm derived from ectopic testicular xenografts. *Reproduction.* 139: 331-335.
- Oktay, K., Newton, H., Mullan, J. and Gosden, R.G. 1998. Development of human primordial follicles to antral stages in SCID/hpg mice stimulated with follicle stimulating hormone. *Hum Reprod.* 13: 1133-1138.
- Senbon, S., Ota, A., Tachibana, M. and Miyano, T. 2003. Bovine oocytes in secondary follicles grow and acquire meiotic competence in severe combined immunodeficient mice. *Zygote.* 11: 139-149.
- Schlatt, S., Kim, S.S. and Gosden, R. 2002. Spermatogenesis and steroidogenesis in mouse, hamster and monkey testicular tissue after cryopreservation and heterotopic grafting to castrated hosts. *Reproduction.* 124: 339-346.
- Snedaker, A.K., Honaramooz, A. and Dobrinski, I. A game of cat and mouse: xenografting of testis tissue from domestic kittens results in complete cat spermatogenesis in a mouse host. *J Androl.* 2004; 25: 926-930.
- Snow, M., Cox, S.L., Jenkin, G., Trounson, A. and Shaw, J. Generation of live young from xenografted mouse ovaries. *Science.* 297: 2227.
- Somfai, T., Ozawa, M., Noguchi, J., Kaneko, H., Nakai, M., Maedomari, N., Ito, J., Kashiwazaki, N., Nagai, T. and Kikuchi, K. 2009. Live piglets derived from *in vitro*-produced zygotes vitrified at the pronuclear stage. *Biol Reprod.* 80: 42-49.
- Somfai, T., Kikuchi, K. and Nagai, T. 2012. Factors affecting cryopreservation of porcine oocytes. *J Reprod Dev.* 58: 17-24.
- Somfai, T., Yoshioka, K., Tanihara, F., Kaneko, H., Noguchi, J., Kashiwazaki, N., Nagai, T. and Kikuchi, K. 2014. Generation of live piglets from cryopreserved oocytes for the first time using a defined system for *in vitro* embryo production. *PLoSOne.* 9: e97731.
- Weissman, A., Gotlieb, L., Colgan, T., Jurisicova, A., Greenblatt, E.M. and Casper, R.F. 1999. Preliminary experience with subcutaneous human ovarian cortex transplantation in the NODSCID mouse. *Biol Reprod.* 60: 1462-1467.
- Zeng, W., Avelar, G.F., Rathi, R., Franca, L.R. and Dobrinski, I. 2006. The length of the spermatozoatogenic cycle is conserved in porcine and ovine testis xenografts. *J Androl.* 27: 527-533.
- Zeng, W., Rathi, R., Pan, H. and Dobrinski, I. 2007. Comparison of global gene expression between porcine testis tissue xenografts and porcine testis *in situ*. *Mol Reprod Dev.* 74: 674-679.